KEEP OUT OF REACH OF CHILDREN WARNING



GREENGUARD™

PROTECTANT AND SYSTEMIC **FUNGICIDE**

Controls fungal diseases on garden plants



Eye corrosive



Ecotoxic



Manager (Approvals Operations) Warce Zinzley



Greenguard™

CONTENTS MAKE 5-40 LITRES OF SPRAY

IMPORTANT: PLEASE READ THE LABEL COMPLETELY BEFORE USE

methyl in the form of a suspension concentrate ACTIVE INGREDIENTS: Contains 102g/litre chlorothalonil and 125g/litre thiophanate-

if taken orally at repeated high doses. Suspected of causing cancer or genetic defects. May **WARNING:** This product is corrosive and may cause severe eye irritation and eye damage. May be harmful if swallowed, inhaled or absorbed through the skin. Toxic to human kidneys Toxic to the soil environment, mammals, birds, and fish. cause mild skin irritation or an allergic skin reaction. Very toxic to the aquatic environment

NOT allow spray to drift onto desirable plants, aquatic environment or outside the target DO NOT contaminate any water supply or fishponds with product or empty container. DO inhalation of spray mist. Wear protective gloves and eye protection when mixing or using PRECAUTIONS: Keep out of reach of children. Avoid contact with the eyes, skin and Use product in well-ventilated areas. Wash hands and exposed skin thoroughly after use Keep children, pets, wildlife and birds away from treated areas until the spray is dry.

within 24 hours RECORD KEEPING: Records of use must be kept if using more than 200mL of product

an approved landfill. Rinse empty container, puncture, and recycle or dispose of safely with domestic rubbish. children and foodstuffs. Dispose of this product only by using according to the label or at DISPOSAL: Store in original container, tightly closed, locked away from

FOR EMERGENCIES ONLY, CALL 0800 734 607 (24 Hours)

respiration if breathing has stopped. splashed in eyes, wash out immediately with running water for at least 15 minutes and seek contact occurs, remove contaminated clothing and flush skin and hair with running water. FIRST AID: If swallowed, do NOT induce vomiting. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If skin or hair If inhaled move the victim to fresh air immediately. Begin artificial

curative and systemic activity. GENERAL INFORMATION: Greenguard is a general garden fungicide with protectant

DIRECTIONS: Spray plants thoroughly Shake well. Measure the correct amount of Greenguard into water and stir Apply at first sign of disease and then at 10-14 day intervals

well. Oplay plants the	well. Oplay plants indivigily. Apply at hist sign of disease and then at 10-14 day intervals.	ally fileli at 10-14 day illelyals.
DISEASE		RATE
ORNAMENTALS	Powdery Mildew, Blackspot,	
(including roses)	ertain	5ml/litre
	leaf spot fungi	
VEGETABLES	Tomatoes: Botrytis, Late blight,	5ml/litre
IN	Sclerotinia	5ml/litre
	Potatoes: Late blight	5ml/litre PRIMAR
	Cabbages: Clubroot	10ml/litre
	Beans: Botrytis, Sclerotinia	RY
		いるがではない。

MINIS

2 0 NOV

(text appearing above this line does not form part of the label) Label date: 07 November 2013

Page 3 of 3

	tilleau, Golden bracelet disease	
	throad Coldon broadlet diagons	
	patch, Anthracnose, Leaf spot, Red	
	(Sclerotinia), Rhizoctonia, Brown	COTULA
30-40ml/litre per 10sq. metres.	Fusarium patch, Dollar spot	TURF, LAWNS,
	flowering)	
	Stonefruit: Brown rot (apply at	
5ml/litre	Blackspot, Botrytis	
5ml/litre	Grapes: Powdery mildew,	FRUIT

It is an offence to use this product on animals

the use of this product must be observed. DO NOT graze treated crops. DO NOT feed any grazing stock to give unacceptable residues in meat and milk, the following restrictions on or other minimum tillage techniques) before sowing pasture or any animal feed crop. part of a treated crop to stock. CULTIVATE treated areas thoroughly (i.e. not direct drilling Yates Greenguard contains minute quantities of HCB, a compound that can be taken up by

the flower. CAUTION: Spraying when flower buds area showing petal colour may result in damage q

WITHHOLDING PERIODS: Beans and potatoes – allow 14 days between spraying and harvest. Grapes – allow 28 days. Stonefruit – do not apply after flowering. All other cro Stonefruit - do not apply after flowering. All other crops

concentrate formulations COMPATIBILITY: Do not mix with Lime Sulphur Bordeaux mixtures, oil or emulsifiable

conditions Registered pursuant to the ACVM Act 1997, No. P4469. See www.foodsafety.govt.nz for registration

 \circledR Yates is a registered trade mark. The front panel of this package is owned under copyright laws by Yates New Zealand.

LIABILITY: This product must be used strictly as directed. Yates will not be liable for any loss or damage arising from a failure to strictly follow the directions for use.

Registered to and distributed by:

DuluxGroup (NZ) Pty Ltd The Gate, 373 Neilson Street, Onehunga, Auckland, (09) 636 2800

www.yates.co.nz

200mL NET

Batch no.:

DOM:



Fcotoxic



